
## Ocean Disclosure Project 2016 SKRETT





Fishery name and location	Fishing gear used	Certification or FIP	Category	Environmental notes
Gulf menhaden Gulf of Mexico United States		Not Certified or in a FIP	В	Bycatch is considered low in this fishery.  No direct impact on PET species.  This fish plays an important role in the marine food web and so potential impacts on the wider marine ecosystem must be monitored.  No impact on seabed.
Araucanian herring Chilean Chile		FIP (D)	C	Bycatch is considered low in this fishery.  No direct impact on PET species.  This fish plays an important role in the marine food web and so potential impacts on the wider marine ecosystem must be monitored.  No impact on seabed.
Lesser sandeel Central Eastern North sea European Union		Not Certified or in a FIP	В	Bycatch is considered low in this fishery.  No direct impact on PET species.  Sandeel is an important prey species for a variety of predators, including fish, marine mammals and seabirds. Fishing for sandeel has been found to affect some surface feeding bird species, especially kittiwake and sandwich tern.  Bottom trawls impact on the seabed.
Anchoveta Chilean regions III and IV (2) Chile		Not Certified or in a FIP	B	Bycatch is considered low in this fishery.  No direct impact on PET species.  This fish plays an important role in the marine food web and so potential impacts on the wider marine ecosystem must be monitored.  No impact on seabed.  *Please refer to the FishSource website for further information: www.fishsource.org

Fishery name and location	Fishing gear used	Certification or FIP	Category	Environmental notes
Anchoveta  Peruvian Northern Central stock Peru		Not Certified or in a FIP	В	Bycatch is considered low in this fishery.  No direct impact on PET species.  This fish plays an important role in the marine food web and so potential impacts on the wider marine ecosystem must be monitored.  No impact on seabed.
Capelin Icelandic Iceland		Not Certified or in a FIP	В	Bycatch is considered low in this fishery.  No direct impact on PET species.  This fish plays an important role in the marine food web and so potential impacts on the wider marine ecosystem must be monitored.  No impact on seabed.
Boarfish North-East Atlantic (2) European Union		Not Certified or in a FIP	B	Bycatch is considered low in this fishery.  No direct impact on PET species.  This fish plays an important role in the marine food web and so potential impacts on the wider marine ecosystem must be monitored.  No impact on seabed.
Lesser sandeel Central Western North Sea (2) European Union		Not Certified or in a FIP	В	Bycatch is considered low in this fishery.  No direct impact on PET species.  This fish plays an important role in the marine food web and so potential impacts on the wider marine ecosystem must be monitored.  Bottom trawls impact on the seabed.

Fishery name and location	Fishing gear used	Certification or FIP	Category	Environmental notes
Lesser sandeel  Shetland (2) European Union		Not Certified or in a FIP	В	Bycatch is considered low in this fishery.  No direct impact on PET species.  This fish plays an important role in the marine food web and so potential impacts on the wider marine ecosystem must be monitored.  Bottom trawls impact on the seabed.
Blue whiting North-East Atlantic EU, Iceland, Faroes, Norway		Not Certified or in a FIP	C	Bycatch is considered low in this fishery.  No direct impact on PET species.  Sandeel is an important prey species for a variety of predators, including fish, marine mammals and seabirds. Fishing for sandeel has been found to affect some surface feeding bird species, especially kittiwake and sandwich tern.  Bottom trawls impact on the seabed.
Norway pout  North Sea  European Union		Not Certified or in a FIP	B	Bycatch is considered low in this fishery.  No direct impact on PET species.  Sandeel is an important prey species for a variety of predators, including fish, marine mammals and seabirds. Fishing for sandeel has been found to affect some surface feeding bird species, especially kittiwake and sandwich tern.  Bottom trawls impact on the seabed.
European pilchard North-West Africa Southern stock Morocco, Mauretania, Senegal		Not Certified or in a FIP	В	Bycatch is considered low in this fishery.  No direct impact on PET species.  Sandeel is an important prey species for a variety of predators, including fish, marine mammals and seabirds. Fishing for sandeel has been found to affect some surface feeding bird species, especially kittiwake and sandwich tern.  Bottom trawls impact on the seabed.

Fishery name and location	Fishing gear used	Certification or FIP	Category	Environmental notes
Lesser sandeel  SE North Sea European Union		Not Certified or in a FIP	C	Bycatch is considered low in this fishery.  No direct impact on PET species.  Sandeel is an important prey species for a variety of predators, including fish, marine mammals and seabirds. Fishing for sandeel has been found to affect some surface feeding bird species, especially kittiwake and sandwich tern.  Bottom trawls impact on the seabed.
Lesser sandeel  Dogger Bank area  European Union		Not Certified or in a FIP	В	Bycatch is considered low in this fishery.  No direct impact on PET species.  Sandeel is an important prey species for a variety of predators, including fish, marine mammals and seabirds. Fishing for sandeel has been found to affect some surface feeding bird species, especially kittiwake and sandwich tern.  Bottom trawls impact on the seabed.
Anchoveta Chilean regions xv-i-ii / Southern Peruvian stock Chile/Peru		Not Certified or in a FIP	C	Bycatch is considered low in this fishery.  No direct impact on PET species.  This fish plays an important role in the marine food web and so potential impacts on the wider marine ecosystem must be monitored.  No impact on seabed.
European pilchard  North-West Africa central stock  Morocco, Mauretania, Senegal		Not Certified or in a FIP	B	Bycatch is considered low in this fishery.  No direct impact on PET species.  This fish plays an important role in the marine food web and so potential impacts on the wider marine ecosystem must be monitored.  No impact on seabed.  *Please refer to the FishSource website for further information: www.fishsource.org













Fishery name and location	Fishing gear used	Certification or FIP	Category	Environmental notes
Capelin  Barents Sea  Russia/Norway		Not Certified or in a FIP	C	Bycatch is considered low in this fishery.  No direct impact on PET species.  This fish plays an important role in the marine food web and so potential impact on the wider marine ecosystem must be monitored.  No impact on seabed.
Anchoveta Chilean regions V-X Chile		Not Certified or in a FIP	C	Bycatch is considered low in this fishery.  No direct impact on PET species.  This fish plays an important role in the marine food web and so potential impact on the wider marine ecosystem must be monitored.  No impact on seabed.
Lesser sandeel  Kattegat (2)  European Union		Not Certified or in a FIP	C	Bycatch is considered low in this fishery.  No direct impact on PET species.  Sandeel is an important prey species for a variety of predators, including fish, marine mammals and seabirds. Fishing for sandeel has been found to affect some surface feeding bird species, especially kittiwake and sandwich tern.  Bottom trawls impact on the seabed.
Lesser sandeel Viking and Bergen Banks ( European Union	2)	Not Certified or in a FIP	C	Bycatch is considered low in this fishery.  No direct impact on PET species.  Sandeel is an important prey species for a variety of predators, including fish, marine mammals and seabirds. Fishing for sandeel has been found to affect some surface feeding bird species, especially kittiwake and sandwich tern.  Bottom trawls impact on the seabed.

NOTE: ALL DATA FROM SKRETTING ANNUAL SUSTAINABILITY REPORT 2013

NOTE: SPECIFIC STOCKS CANNOT BE IDENTIFIED FOR SAND EEL AND PERUVIAN ANCHOVY































